


10/590,871

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification⁶ : C07D 207/00, 401/00, C07K 5/00	A2	(11) International Publication Number: WO 98/19998 (43) International Publication Date: 14 May 1998 (14.05.98)
(21) International Application Number: PCT/EP97/06125 (22) International Filing Date: 5 November 1997 (05.11.97) (30) Priority Data: 08/746,295 7 November 1996 (07.11.96) US (71) Applicant (for all designated States except US): NOVARTIS AG [CH/CH]; Schwarzwaldallee 215, CH-4058 Basel (CH). (72) Inventor; and (75) Inventor/Applicant (for US only): VILLHAUER, Edwin, Bernard [US/US]; 20 Dorothy Drive, Morristown, NJ 07960 (US). (74) Agent: ROTH, Bernhard, M.; Novartis AG, Patent- und Markenabteilung, Lichtstrasse 35, CH-4002 Basel (CH).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: N-SUBSTITUTED 2-CYANOPYRROLIDINES (57) Abstract N-(N'-substituted glycy)-2-cyanopyrrolidines, e.g. the compounds of formula (I) wherein R has various significances, are novel. They inhibit DPP-IV (dipeptidyl-peptidase-IV) activity. They are therefore indicated for use as pharmaceuticals in inhibiting DPP-IV and in the treatment of conditions mediated by DPP-IV, such as non-insulin-dependent diabetes mellitus, arthritis, obesity, osteoporosis and further conditions of impaired glucose tolerance.		



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

N-SUBSTITUTED 2-CYANOPYRROLIDINES

Field

The present invention relates to N-substituted 2-cyanopyrrolidines. More particularly, it provides novel N-glycyl-2-cyanopyrrolidine derivatives.

Background

Dipeptidyl peptidase-IV (**DPP-IV**) is a serine protease which cleaves N-terminal dipeptides from a peptide chain containing, preferably, a proline residue in the penultimate position. Although the biological role of DPP-IV in mammalian systems has not been completely established, it is believed to play an important role in neuropeptide metabolism, T-cell activation, attachment of cancer cells to the endothelium and the entry of HIV into lymphoid cells. DPP-IV is responsible for inactivating glucagon-like peptide-1 (**GLP-1**). More particularly, DPP-IV cleaves the amino-terminal His-Ala dipeptide of GLP-1, generating a GLP-1 receptor antagonist, and thereby shortens the physiological response to GLP-1. Since the half-life for DPP-IV cleavage is much shorter than the half-life for removal of GLP-1 from circulation, a significant increase in GLP-1 bioactivity (5- to 10-fold) is anticipated from DPP-IV inhibition. Since GLP-1 is a major stimulator of pancreatic insulin secretion and has direct beneficial effects on glucose disposal, DPP-IV inhibition appears to represent an attractive approach for treating non-insulin-dependent diabetes mellitus (**NIDDM**).


Although a number of DPP-IV inhibitors have been described, all have limitations relating to potency, stability or toxicity. Accordingly, a great need exists for novel DPP-IV inhibitors which are useful in treating conditions mediated by DPP-IV inhibition and which do not suffer from the above-mentioned limitations.

Summary of the invention

The invention provides novel N-(N'-substituted glycyI)-2-cyanopyrrolidines which are effective as DPP-IV inhibitors in treating conditions mediated by DPP-IV. It also concerns corresponding pharmaceutical compositions, a process for their preparation, a method of inhibiting DPP-IV comprising administering to a patient in need of such treatment a therapeutically effective amount thereof, the compounds for use as a pharmaceutical, and their use in a process for the preparation of a medicament for treating a condition mediated by DPP-IV.

Detailed description

The invention concerns N-(N'-substituted glycyI)-2-cyanopyrrolidines, hereinafter briefly named "**the compounds of the invention**"; more particularly, it concerns compounds of formula I:



wherein **R** is:

a) $R_1R_{1a}N(CH_2)_m$ wherein

R_1 is a pyridinyl or pyrimidinyl moiety optionally mono- or independently disubstituted with (C_{1-4}) alkyl, (C_{1-4}) alkoxy, halogen, trifluoromethyl, cyano or nitro; or phenyl optionally mono- or independently disubstituted with (C_{1-4}) alkyl, (C_{1-4}) alkoxy or halogen;

R_{1a} is hydrogen or (C_{1-8}) alkyl; and

m is 2 or 3;

b) (C_{3-12}) cycloalkyl optionally monosubstituted in the 1-position with (C_{1-3}) hydroxyalkyl;

c) $R_2(CH_2)_n$ wherein either

R_2 is phenyl optionally mono- or independently di- or independently trisubstituted with (C_{1-4}) alkyl, (C_{1-4}) alkoxy, halogen or phenylthio optionally monosubstituted in the phenyl ring with hydroxymethyl; or is (C_{1-8}) alkyl; a [3.1.1]bicyclic carbocyclic moiety optionally mono- or plurisubstituted with (C_{1-8}) alkyl;

- a pyridinyl or naphthyl moiety optionally mono- or independently disubstituted with (C₁₋₄)alkyl, (C₁₋₄)alkoxy or halogen; cyclohexene; or adamantyl; and
n is 1 to 3; or
R₂ is phenoxy optionally mono- or independently disubstituted with (C₁₋₄)alkyl, (C₁₋₄)alkoxy or halogen; and
n is 2 or 3;
- d) (R₃)₂CH(CH₂)₂- wherein each R₃ independently is phenyl optionally mono- or independently disubstituted with (C₁₋₄)alkyl, (C₁₋₄)alkoxy or halogen;
- e) R₄(CH₂)_p- wherein R₄ is 2-oxopyrrolidinyl or (C₂₋₄)alkoxy and
p is 2 to 4;
- f) isopropyl optionally monosubstituted in 1-position with (C₁₋₃)hydroxyalkyl;
- g) R₅ wherein R₅ is: indanyl; a pyrrolidinyl or piperidinyl moiety optionally substituted with benzyl; a [2.2.1]- or [3.1.1]bicyclic carbocyclic moiety optionally mono- or plurisubstituted with (C₁₋₈)alkyl; adamantyl; or (C₁₋₈)alkyl optionally mono- or independently plurisubstituted with hydroxy, hydroxymethyl or phenyl optionally mono- or independently disubstituted with (C₁₋₄)alkyl, (C₁₋₄)alkoxy or halogen;
- in free form or in acid addition salt form.

The compounds of formula I can exist in free form or in acid addition salt form. Salt forms may be recovered from the free form in known manner and vice-versa. Acid addition salts may e.g. be those of pharmaceutically acceptable organic or inorganic acids. Although the preferred acid addition salts are the hydrochlorides, salts of methanesulfonic, sulfuric, phosphoric, citric, lactic and acetic acid may also be utilized.

The compounds of the invention may exist in the form of optically active isomers or diastereoisomers and can be separated and recovered by conventional techniques, such as chromatography.

"Alkyl" and "alkoxy" are either straight or branched chain, of which examples of the latter are isopropyl and tert-butyl.

R preferably is a), b) or e) as defined above. R₁ preferably is a pyridinyl or pyrimidinyl moiety optionally substituted as defined above. R_{1a} preferably is hydrogen. R₂ preferably is phenyl optionally substituted as defined above. R₃ preferably is unsubstituted phenyl.

R₄ preferably is alkoxy as defined above. R₅ preferably is optionally substituted alkyl as defined above. m preferably is 2. n preferably is 1 or 2, especially 2. p preferably is 2 or 3, especially 3.

Pyridinyl preferably is pyridin-2-yl; it preferably is unsubstituted or monosubstituted, preferably in 5-position. Pyrimidinyl preferably is pyrimidin-2-yl. It preferably is unsubstituted or monosubstituted, preferably in 4-position. Preferred as substituents for pyridinyl and pyrimidinyl are halogen, cyano and nitro, especially chlorine.

When it is substituted, phenyl preferably is monosubstituted; it preferably is substituted with halogen, preferably chlorine, or methoxy. It preferably is substituted in 2-, 4- and/or 5-position, especially in 4-position.

(C₃₋₁₂)cycloalkyl preferably is cyclopentyl or cyclohexyl. When it is substituted, it preferably is substituted with hydroxymethyl. (C₁₋₄)alkoxy preferably is of 1 or 2 carbon atoms, it especially is methoxy. (C₂₋₄)alkoxy preferably is of 3 carbon atoms, it especially is isopropoxy. Halogen is fluorine, chlorine, bromine or iodine, preferably fluorine, chlorine or bromine, especially chlorine. (C₁₋₈)alkyl preferably is of 1 to 6, preferably 1 to 4 or 3 to 5, especially of 2 or 3 carbon atoms, or methyl. (C₁₋₄) alkyl preferably is methyl or ethyl, especially methyl. (C₁₋₃)hydroxyalkyl preferably is hydroxymethyl.

A [3.1.1]bicyclic carbocyclic moiety optionally substituted as defined above preferably is bicyclo[3.1.1]hept-2-yl optionally disubstituted in 6-position with methyl, or bicyclo[3.1.1]hept-3-yl optionally trisubstituted with one methyl in 2-position and two methyl groups in 6-position. A [2.2.1]bicyclic carbocyclic moiety optionally substituted as defined above preferably is bicyclo[2.2.1]hept-2-yl.

Naphthyl preferably is 1-naphthyl. Cyclohexene preferably is cyclohex-1-en-1-yl. Adamantyl preferably is 1- or 2-adamantyl.

A pyrrolidinyl or piperidinyl moiety optionally substituted as defined above preferably is pyrrolidin-3-yl or piperidin-4-yl. When it is substituted it preferably is N-substituted.

A preferred group of compounds of the invention is the compounds of formula I wherein R is **R'** (**compounds Ia**), whereby R' is:

- $R_1'NH(CH_2)_2$ - wherein R_1' is pyridinyl optionally mono- or independently disubstituted with halogen, trifluoromethyl, cyano or nitro; or unsubstituted pyrimidinyl;
 - (C_{3-7}) cycloalkyl optionally monosubstituted in 1-position with (C_{1-3}) hydroxyalkyl;
 - $R_4'(CH_2)_3$ - wherein R_4' is (C_{2-4}) alkoxy; or
 - R_5 , wherein R_5 is as defined above;
- in free form or in acid addition salt form.

More preferred compounds of the invention are those compounds of formula I wherein R is **R''** (**compounds Ib**), whereby R'' is:

- $R_1''NH(CH_2)_2$ - wherein R_1'' is pyridinyl mono- or independently disubstituted with halogen, trifluoromethyl, cyano or nitro;
 - (C_{4-6}) cycloalkyl monosubstituted in 1-position with (C_{1-3}) hydroxyalkyl;
 - $R_4'(CH_2)_3$ - wherein R_4' is as defined above; or
 - R_5' wherein R_5' is a [2.2.1]- or [3.1.1]bicyclic carbocyclic moiety optionally mono- or plurisubstituted with (C_{1-8}) alkyl; or adamantyl;
- in free form or in acid addition salt form.

Even more preferred compounds of the invention are the compounds of formula I wherein R is **R'''** (**compounds Ic**), whereby R''' is:

- $R_1''NH(CH_2)_2$ - wherein R_1'' is as defined above;
 - (C_{4-6}) cycloalkyl monosubstituted in 1-position with hydroxymethyl;
 - $R_4'(CH_2)_3$ - wherein R_4' is as defined above; or
 - R_5'' wherein R_5'' is adamantyl;
- in free form or in acid addition salt form.

A further group of compounds of the invention is **compounds Ip**, wherein R is **R^P**, which is:

- a) $R_1^P NH(CH_2)_2$ - wherein R_1^P is a pyridinyl or pyrimidinyl moiety optionally mono- or independently disubstituted with halogen, trifluoromethyl, cyano or nitro;
- b) (C_{3-7}) cycloalkyl optionally monosubstituted in 1-position with (C_{1-3}) hydroxyalkyl;
- c) $R_2^P(CH_2)_2$ - wherein R_2^P is phenyl optionally mono- or independently di- or independently trisubstituted with halogen or (C_{1-3}) alkoxy;
- d) $(R_3^P)_2CH(CH_2)_2$ - wherein each R_3^P independently is phenyl optionally monosubstituted with halogen or (C_{1-3}) alkoxy;
- e) $R_4(CH_2)_3$ - wherein R_4 is as defined above; or
- f) isopropyl optionally monosubstituted in 1-position with (C_{1-3}) hydroxyalkyl; in free form or in pharmaceutically acceptable acid addition salt form.


A further group of compounds of the invention is **compounds Is**, wherein R is **R^S**, which is:

- a) $R_1^S R_{1a}^S (CH_2)_{ms}$ - wherein R_1^S is pyridinyl optionally mono- or independently disubstituted with chlorine, trifluoromethyl, cyano or nitro; pyrimidinyl optionally monosubstituted with chlorine or trifluoromethyl; or phenyl;
 R_{1a}^S is hydrogen or methyl; and
 ms is 2 or 3;
- b) (C_{3-12}) cycloalkyl optionally monosubstituted in 1-position with hydroxymethyl;
- c) $R_2^S(CH_2)_{ns}$ - wherein either
 R_2^S is phenyl optionally mono- or independently di- or independently trisubstituted with halogen, alkoxy of 1 or 2 carbon atoms or phenylthio monosubstituted in the phenyl ring with hydroxymethyl;
 (C_{1-6}) alkyl; 6,6-dimethylbicyclo[3.1.1]hept-2-yl; pyridinyl;
naphthyl; cyclohexene; or adamantyl; and ns is 1 to 3; or
 R_2^S is phenoxy; and ns is 2;
- d) (3,3-diphenyl)propyl;

-7-

- e) $R_4^s(CH_2)_{ps}$ wherein R_4^s is 2-oxopyrrolidin-1-yl or isopropoxy and ps is 2 or 3;
- f) isopropyl optionally monosubstituted in 1-position with hydroxymethyl;
- g) R_5^s wherein R_5^s is: indanyl; a pyrrolidinyl or piperidinyl moiety optionally N-substituted with benzyl; bicyclo[2.2.1]hept-2-yl; 2,6,6-trimethylbicyclo[3.1.1]hept-3-yl; adamantyl; or (C_{1-8}) alkyl optionally mono- or independently disubstituted with hydroxy, hydroxymethyl or phenyl; in free form or in acid addition salt form.

The compounds of the invention may be prepared by a process which comprises coupling a reactive (2-cyanopyrrolidino)carbonylmethylene compound with an appropriate substituted amine; more particularly, for the preparation of the compounds of formula I it comprises reacting a compound of formula II



wherein X is a reactive group,
with a compound of formula III



wherein R is as defined above,
and recovering the resultant compound of formula I in free form or in acid addition salt form.

X preferably is a halogen such as bromine, chlorine or iodine.


The process of the invention may be effected in conventional manner.

The compound of formula II is preferably reacted with at least 3 equivalents of a primary amine of formula III. The reaction is conveniently conducted in the presence of an inert, organic solvent, preferably a cyclic ether such as tetrahydrofuran. The temperature preferably is of from about 0° to about 35°C, preferably between about 0° and about 25°C.

The compounds of the invention may be isolated from the reaction mixture and purified in conventional manner, e.g. by chromatography.

The starting materials may also be prepared in conventional manner.

The compounds of formula II may e.g. be prepared by the following two-step reaction scheme:



Step 1 involves the reaction of the pyrrolidine of formula IV with a slight molar excess of a haloacetylhalide such as bromoacetyl bromide or chloroacetyl chloride and triethylamine and a catalytic amount of dimethylaminopyridine (DMAP). The reaction conveniently is conducted in the presence of an inert, organic solvent, preferably a chlorinated, aliphatic hydrocarbon such as methylene chloride, at a temperature of from about 0° to about 25°C, preferably at a temperature between about 0° and about 15°C.

Step 2 concerns the dehydration of the compound of formula V, prepared in Step 1, with at least 2 equivalents of trifluoroacetic anhydride (TFAA). The dehydration preferably is conducted in the presence of an inert, organic solvent such as tetrahydrofuran or a chlorinated, aliphatic hydrocarbon such as methylene chloride, at a temperature of from about 0° to about 25°C, preferably at a temperature between about 0° and about 15°C.

Insofar as its preparation is not particularly described herein, a compound used as starting material is known or may be prepared from known compounds in known manner or analogously to known methods or analogously to methods described in the Examples.

The following Examples illustrate the invention. All temperatures are in degrees Celsius.

Example 1: 1-[2-[(5-Chloropyridin-2-yl)amino]ethylamino]acetyl-2-cyano-(S)-pyrrolidine

To a 500 ml flask is added 16.6 g of 2-[(5-chloropyridin-2-yl)amino]ethylamine and 100 ml of tetrahydrofuran and the mixture is cooled in an ice bath. To the cooled mixture is added 7.0 g of (2-cyanopyrrolidino)carbonylmethylene-(S)-bromide dissolved in 30 ml of tetrahydrofuran. The resultant mixture is stirred for 2 hours at 0°, the solvent is removed by rotovaping and the mixture is partitioned between ethyl acetate and water. The product is then extracted into the ethyl acetate layer and the aqueous layer is then washed twice with ethyl acetate. The combined organic layers are washed successively with water and brine, dried over sodium sulfate and concentrated to obtain the desired free base compound in crude form. The crude form is then purified on silica gel employing a mixture of 5% methanol in methylene chloride as the eluent to yield the **title compound in free base form** as a light brown oil.

After dissolving the **free base** in 30 ml of dry tetrahydrofuran, **hydrogen chloride** gas is bubbled into the solution for five seconds. The off-white precipitate that forms is filtered, washed with dry tetrahydrofuran and the solvent is removed by high vacuum pumping to obtain the **title compound in dihydrochloride acid addition salt form** (off-white solid; m.p. 265°-267°; NMR: see * at bottom of Table hereunder).

The starting material is obtained as follows:

a) 22.37 g of (S)-2-carbamoylpyrrolidine, 30.1 ml of triethylamine and 30.0 mg of dimethylaminopyridine (DMAP) are dissolved in 200 ml of methylene chloride and the solution is then added, dropwise, to an ice-cold solution of 18.8 ml of bromoacetyl bromide in 192 ml of methylene chloride, over a period of 60 minutes under a calcium sulfate drying tube. The resultant solution is stirred for 2 hours at ice-water temperature under a calcium sulfate drying tube, then poured into 3.5 liters of ethyl acetate. The resultant precipitate is filtered, washed with ethyl acetate, and the filtrate is concentrated to obtain (2-carbamoylpyrrolidino)-carbonylmethylene-(S)-bromide (hard yellow taffy).

-10-

b) 50.0 g of the **bromide compound** prepared in a) above is dissolved in 300 ml of methylene chloride and the solution is cooled in an ice water bath under a calcium sulfate drying tube. The cooled solution is then poured into 60.2 ml of **trifluoroacetic anhydride** over a 2 minute period, the resultant solution is stirred at ice-water temperature under a calcium sulfate drying tube for 4 hours, and partitioned between methylene chloride and saturated aqueous sodium bicarbonate. The product is extracted into the methylene chloride layer and the aqueous layer is washed twice with methylene chloride. The combined organic layers are washed successively with water and brine and then dried over sodium sulfate. The solution is filtered and the solvent is removed by rotovaping and high vacuum pumping to obtain **(2-cyanopyrrolidino)carbonylmethylene-(S)-bromide** (dark yellow solid).

The following compounds of the invention, of formula I, are obtained in analogous manner by reacting a corresponding compound of formula II with a corresponding compound of formula III (in the following Table, where only an acid addition salt form of a compound of the invention is mentioned, the compound is recovered from the free base without isolation thereof):

Example No.	R	Form	Analogously to Ex. No.	Characterization data
2	2-[(5-CF ₃ -pyridin-2-yl)amino]ethyl	b	1 ¹⁾	golden oil; NMR*
3	2-[(5-cyanopyridin-2-yl)amino]ethyl	b	1	golden oil
		dch	1	off-white precipitate, m.p. 155-157°; NMR*; [α] _D ²⁰ = - 77.2° (c=0.012, MeOH)
4	2-[(pyrimidin-2-yl)amino]ethyl	b	2 ^{1a)}	golden oil; NMR*
5	(1-hydroxymethyl)cyclopent-1-yl	b	1 ²⁾	yellow solid; m.p. 65-67°; NMR*
6	2-[(pyridin-2-yl)amino]ethyl	b	2 ³⁾	golden oil; NMR*
7A	2-[(4-chloropyrimidin-2-yl)amino]ethyl	b	2	tan solid; NMR*
7B	2-[(3-chloropyridin-2-yl)amino]ethyl	b	2	golden oil; NMR*
7C	2-[(4-CF ₃ -pyrimidin-2-yl)amino]ethyl	b	2	golden oil; NMR*
7D	2-(2-chlorophenyl)ethyl	b	2	NMR*
7E	(3,3-diphenyl)propyl	b	2	NMR*
8	2-[(5-nitropyridin-2-yl)amino]ethyl	b	5 ^{3a)}	bright yellow thick oil; NMR*
9A	2-[(3-chloro-5-CF ₃ -pyridin-2-yl)amino]ethyl	b	2 ^{3b)}	golden oil; NMR*
9B	2-[(3-CF ₃ -pyridin-2-yl)amino]ethyl	b	2 ^{3b)}	golden oil; NMR*
9C	2-[(3,5-dichloropyridin-2-yl)amino]ethyl	b	2 ^{3b)}	golden oil; NMR*
10	cyclopent-1-yl	b	2	tan solid
		ch	1	white solid; NMR*
11	2-(2-bromo-4,5-dimethoxyphenyl)ethyl	b	5 ^{3a)}	clear light yellow, thick oil; NMR*
12	3-(isopropoxy)propyl	b	1	brown oil
		ch	1	white solid, m.p. 174-176°; NMR*

Example No.	R	Form	Analogously to Ex. No.	Characterization data
13	2-hydroxy-1,1-dimethylethyl	b	2 ⁴⁾	golden oil
14	3-(2-oxopyrrolidin-1-yl)propyl	ch	1	brown solid; NMR*
15	1-hydroxymethylcyclohexyl	b	2 ⁵⁾	golden oil
16	2-(4-ethoxyphenyl)ethyl	ch	1	tan solid; NMR*
17	1-phenylmethyl-3-(R)-pyrrolidinyl	b	1 ⁶⁾	yellow waxy solid; m.p. 93°; ¹³ C-NMR: 118.1 (ppm)
18	2-(4-methoxyphenyl)ethyl	ch	1	white solid; m.p. 182-184°; ¹³ C-NMR: 121.4 (ppm)
19	2-(3-methoxyphenyl)ethyl	dch	1 ⁷⁾	off-white solid; m.p. 175-177°; ¹³ C-NMR: 121.5 (ppm)
20	(1-naphthalenyl)methyl	ch	1	white solid; m.p. 185-187°; ¹³ C-NMR: 121.4 (ppm)
21	3-phenylpropyl	ch	1	light yellow solid; m.p. 172-174°; ¹³ C-NMR: 119.25 (ppm)
22	3-[(phenyl)(methyl)amino]propyl	ch	1	light yellow solid; m.p. 130-135°; ¹³ C-NMR: 119.29 (ppm)
23	2-(3,4-dimethoxyphenyl)ethyl	ch	1	off-white fluffy solid; ¹³ C-NMR: 119.26 (ppm)
24	cycloheptyl	dch	1	white solid; m.p. 96-98° (foams); ¹³ C-NMR: 121.6 (ppm)
25	[1S[1α,2α(S*),5α]]-(6,6-dimethylbicyclo[3.1.1]-hept-2-yl)methyl	ch	1	white solid; m.p. 170-172°; ¹³ C-NMR: 121.5 (ppm)
26	2-(2,5-dimethoxyphenyl)ethyl	ch	1	white solid; m.p. 68-70°; ¹³ C-NMR: 121.4 (ppm)
27	2-(1-cyclohexen-1-yl)ethyl	ch	1 ⁸⁾	white solid; m.p. 275-279° (dec.); ¹³ C-NMR: 119.17 (ppm)
28	cyclohexyl	ch	1	white fluffy solid; m.p. 65-67°; ¹³ C-NMR: 119.25 (ppm)
29	[1S[1α,2α(S*),5α]]-bicyclo[2.2.1]hept-2-yl	ch	1	off-white fluffy solid; m.p. 162-164°; ¹³ C-NMR: 119.27 (ppm)
30	2-(2-pyridinyl)ethyl	ch	1	white fluffy solid; m.p. 182-184°; ¹³ C-NMR: 119.28 (ppm)
31	(2-phenylamino)ethyl	ch	1 ⁹⁾	white solid; m.p. 98-100°; ¹³ C-NMR: 118.36 (ppm)
32	3,3-dimethylbutyl	dch	1	white solid; m.p. 95-97°; ¹³ C-NMR: 121.5 (ppm)
33	[1S[1α,2β,3α(S*),5α]]-2,6,6-trimethylbicyclo[3.1.1]-hept-3-yl	dch	1	white solid; m.p. 124-126°; ¹³ C-NMR: 121.4 (ppm)
		ch	1	white solid; m.p. 164-166°; ¹³ C-NMR: 121.5 (ppm)
		ch	1 ¹⁰⁾	white solid; m.p. 82-84°; ¹³ C-NMR: 121.5 (ppm)

Example No.	R	Form	Analogously to Ex. No.	Characterization data
34	[S,S]-(1-hydroxymethyl)propyl	ch	1 ¹¹⁾	off-white solid; m.p. 80-82°; ¹³ C-NMR: 118.2 (ppm)
35	[2-[(2-hydroxymethyl)phenyl]thio]phenylmethyl	ch	1 ¹²⁾	yellow solid; m.p. 65-67°; ¹³ C-NMR: 121.4 (ppm)
36	2-(2-methoxyphenyl)ethyl	ch	1	off-white solid; m.p. 174-176°; ¹³ C-NMR: 121.7 (ppm)
37	5-hydroxypentyl	ch	1	sticky light-green solid; ¹³ C-NMR: 121.67 (ppm)
38	cyclobutyl	ch	1	off-white solid; m.p. 274-278° (dec.); ¹³ C-NMR: 121.64 (ppm)
39	2-(2,4-dichlorophenyl)ethyl	ch	1	white fluffy solid; m.p. 154-156°; ¹³ C-NMR: 121.48 (ppm)
40	1-(S)-(+)-hydroxymethyl-3-methylbutyl	ch	1 ¹³⁾	light yellow solid; m.p. 65-66°; ¹³ C-NMR: 117.99 (ppm)
41	[1R*,2S*]-2-hydroxy-2-phenylethyl	ch	1 ¹⁴⁾	light yellow solid; m.p. 82-83°; ¹³ C-NMR: 118.35 (ppm)
42	2-(2-fluorophenyl)ethyl	ch	1	white fluffy solid; m.p. 160-162°; ¹³ C-NMR: 121.70 (ppm)
43	cyclopropyl	ch	1	off-white solid; m.p. 170-172°; ¹³ C-NMR: 121.62 (ppm)
44	[1S(1S,2S,3S,5R)]-2,6,6-trimethylbicyclo-[3.1.1]hept-3-yl	ch	1 ¹⁵⁾	white solid; m.p. 84-86°; ¹³ C-NMR: 121.8 (ppm)
45	(2-phenoxy)ethyl	ch	1	sticky golden solid; ¹³ C-NMR: 121.7 (ppm)
46	2-(3,5-dimethoxyphenyl)ethyl	ch	1	white fluffy solid; m.p. 74-76°; ¹³ C-NMR: 121.66 (ppm)
47	1-adamantyl	ch	1	white solid; m.p. 240-242°; ¹³ C-NMR: 121.80 (ppm)
48	1,1,3,3-tetramethylbutyl	ch	1	white fluffy solid; m.p. 68-70°; ¹³ C-NMR: 121.55 (ppm)
49	2-adamantyl	ch	1	off-white solid; m.p. 122-124°; ¹³ C-NMR: 121.69 (ppm)
50	1,1-dimethylpropyl	ch	1	white fluffy solid; m.p. 62-64°; ¹³ C-NMR: 121.53 (ppm)
51	benzyl	ch	1	white solid; m.p. 58-60°; ¹³ C-NMR: 121.38 (ppm)
52	1,1-dimethylethyl	ch	1	white solid; m.p. 226-228°; ¹³ C-NMR: 121.56 (ppm)
53	(2-adamantyl)methyl	ch	1	white solid; m.p. 158-160°; ¹³ C-NMR: 121.53 (ppm)
54	2-phenylethyl	ch	1	white solid; m.p. 275-280° (dec.); ¹³ C-NMR: 121.52 (ppm)
55	pentyl	ch	1	white solid; m.p. 176-178°; ¹³ C-NMR: 121.67 (ppm)
56	butyl	ch	1	white solid; m.p. 180-182°; ¹³ C-NMR: 121.53 (ppm)

Example No.	R	Form	Analogously to Ex. No.	Characterization data
57	cyclododecyl	ch	1	white fluffy solid; ¹³ C-NMR: 121.52 (ppm)
58	cyclooctyl	ch	1	white fluffy solid; ¹³ C-NMR: 121.64 (ppm)
59	propyl	ch	1	white solid; m.p. 193-194°; ¹³ C-NMR: 121.57 (ppm)
60	ethyl	ch	1	off-white sticky solid; ¹³ C-NMR: 121.67 (ppm)
61	heptyl	ch	1	white solid; m.p. 170-172°; ¹³ C-NMR: 121.7 (ppm)
62	hexyl	ch	1	white solid; m.p. 174-176°; ¹³ C-NMR: 121.75 (ppm)
63	3-[(5-cyano-2-pyridinyl)amino]propyl	dch	1 ¹⁶⁾	white sticky solid; m.p. 210-212°; ¹³ C-NMR: 119.33 (ppm)
64	1-ethylpropyl	ch	1	white fluffy sticky solid; ¹³ C-NMR: 119.35 (ppm)
65	2,3-dihydro-1H-inden-2-yl	ch	1	white solid; m.p. 182-184°; ¹³ C-NMR: 121.38 (ppm)
66	1-benzylpiperidin-4-yl	ch	1	white solid; m.p. 280-283° (dec.); ¹³ C-NMR: 121.39 (ppm)

b = in free base form ch = in monohydrochloride form; dch = in dihydrochloride form

dec. = decomposes; m.p. = melting point

- ¹⁾ starting from 2-[(5-trifluoromethyl)pyridin-2-yl]amino]ethylamine in tetrahydrofuran, using 5 % methanol in methylene chloride as eluent
- ^{1a)} using 10 % methanol in methylene chloride as eluent
- ²⁾ reacting (1-hydroxymethyl)cyclopentylamine in anhydrous tetrahydrofuran for 18 h at room temperature under a calcium sulfate drying tube
- ³⁾ using methylene chloride, methanol and ammonium hydroxide 90:10:0.5 as eluent
- ^{3a)} using flash chromatography, employing a mixture of 5 % methanol in methylene chloride as eluent
- ^{3b)} using 3 % methanol in methylene chloride as eluent
- ⁴⁾ using methylene chloride, methanol and ammonium hydroxide 80:20:1 as eluent
- ⁵⁾ using methylene chloride, methanol and ammonium hydroxide 90:10:1 as eluent
- ⁶⁾ reacting (1-aminocyclohexane)methanol in tetrahydrofuran with 1-chloroacetyl-2-(S)-cyanopyrrolidine (prepared from chloroacetylchloride and L-prolinamide and reaction of the resultant product with trifluoroacetic anhydride) at ice-temperature under a calcium sulfate drying tube, and purifying the reaction mixture on silicagel employing 5 % methanol in methylene chloride as eluent

- ⁷⁾ using (3R)-(-)-1-benzyl-3-aminopyrrolidine as starting material
⁸⁾ using (-)-cis-myrtanylamine as starting material
⁹⁾ using (+/-)-exo-2-aminonorbornane as starting material
¹⁰⁾ using (1R,2R,3R,5S)-(-)-isopinocampheylamine as starting material
¹¹⁾ using (S)-(+)-2-amino-1-butanol as starting material
¹²⁾ using 2-(2-aminomethyl)phenylthio)benzyl alcohol as starting material
¹³⁾ using (S)-(+)-leucinol as starting material
¹⁴⁾ using (1R,2S)-(-)-norephedrine as starting material
¹⁵⁾ using (1S,2S,3S,5R)-(+)-isopinocampheylamine as starting material
¹⁶⁾ using 2-(3-aminopropylamino)-5-cyanopyridine as starting material

* NMR:

Compound #	¹³ C-NMR (MHz, solvent) d ppm (CN)
Ex. 5	¹³ C NMR (75 MHz, CD ₃ OD) d 119.64 ppm (CN)
Ex. 12	¹³ C NMR (75 MHz, D ₂ O) d 121.63 ppm (CN)
Ex. 1	¹³ C NMR (75 MHz, D ₂ O) d 121.60 ppm (CN)
Ex. 3	¹³ C NMR (75 MHz, D ₂ O) d 120.42 ppm (CN)
Ex. 8	¹³ C NMR (75 MHz, DMSO) d 119.13 ppm (CN)
Ex. 7B	¹³ C NMR (75 MHz, CDCl ₃) d 118.23 ppm (CN)
Ex. 9A	¹³ C NMR (75 MHz, CD ₃ OD) d 119.68 ppm (CN)
Ex. 9B	¹³ C NMR (75 MHz, CD ₃ OD) d 119.66 ppm (CN)

Compound #	¹³ C NMR (MHz, solvent) d ppm (CN)
Ex. 9C	¹³ C NMR (75 MHz, CD ₃ OD) d 119.68 ppm (CN)
Ex. 6	¹³ C NMR (75 MHz, CD ₃ OD) d 119.84 ppm (CN)
Ex. 7C	¹³ C NMR (75 MHz, CDCl ₃) d 118.23 ppm (CN)
Ex. 2	¹³ C NMR (75 MHz, CD ₃ OD) d 119.68 ppm (CN)
Ex. 7A	¹³ C NMR (75 MHz, CD ₃ OD) d 119.66 ppm (CN)
Ex. 4	¹³ C NMR (75 MHz, CD ₃ OD) d 119.66 ppm (CN)
Ex. 10	¹³ C NMR (75 MHz, D ₂ O) d 121.69 ppm (CN)
Ex. 11	¹³ C NMR (75 MHz, CDCl ₃) d 118.31 ppm (CN)
Ex. 7D	¹³ C NMR (75 MHz, CD ₃ OD) d 119.63 ppm (CN)
Ex. 7E	¹³ C NMR (75 MHz, CD ₃ OD) d 119.64 ppm (CN)
Ex. 13	¹³ C NMR (75 MHz, D ₂ O) d 121.52 ppm (CN)
Ex. 14	¹³ C NMR (75 MHz, D ₂ O) d 121.52 ppm (CN)

The compounds of the invention in free form or in pharmaceutically acceptable acid addition salt form, hereinafter briefly named "**the agents of the invention**", in particular, the compounds of formula I in free form or in pharmaceutically acceptable acid addition salt form, possess pharmacological activity. They are therefore indicated for use as pharmaceuticals.

In particular, they inhibit DPP-IV. This activity may be demonstrated employing the **Caco-2 DPP-IV assay**, which measures the ability of test compounds to inhibit DPP-IV activity from human colonic carcinoma cell extracts. The human colonic carcinoma cell line Caco-2 can be obtained from the American Type Culture Collection (ATCC HTB 37). Differentiation of the cells to induce DPP-IV expression is accomplished as described by Reisher et al. in Proc.Natl.Acad.Sci.USA **90** (1993) 5757-5761. Cell extract is prepared from cells solubilized in 10 mM Tris-HCl, 0.15 M NaCl, 0.04 t.i.u. (trypsin inhibitor unit) aprotinin, 0.5% non-ionic detergent P40, pH 8.0, which is centrifuged at 35 000 g for 30 min at 4°C to remove cell debris. The assay is conducted by adding 20 mg solubilized Caco-2 protein, diluted to a final volume of 125 ml in assay buffer (25 mM Tris-HCl pH 7.4, 140 mM NaCl, 10 mM KCl, 1% bovine serum albumin) to microtiter plate wells. The reaction is initiated by adding 25ml of 1 mM substrate (H-Alanine-Proline-pNA; pNA is *p*-nitroaniline). The reaction is run at room temperature for 10 minutes after which time a 19 ml volume of 25% glacial acetic acid is added to stop the reaction. Test compounds are typically added as 30 ml additions and the assay buffer volume is reduced to 95 ml. A standard curve of free *p*-nitroaniline is generated using 0-500 mM solutions of free pNA in assay buffer. The curve generated is linear and is used for interpolation of substrate consumption (catalytic activity in nmoles substrate cleaved /min). The endpoint is determined by measuring absorbance at 405 nm in a Molecular Devices UV Max microtiter plate reader. The potency of the test compounds as DPP-IV inhibitors, expressed as IC₅₀, is calculated from 8-point, dose-response curves using a 4-parameter logistic function.

In the above test, IC₅₀ values of from about 10 nM to about 900 nM are obtained with the agents of the invention, e.g. 22 nM for the agent of Example 3.

The DPP-IV inhibition may also be demonstrated by measuring the effects of test compounds on DPP-IV activity in human and rat plasma employing a modified version of the **assay described by Kubota et al.** in Clin.Exp.Immunol. **89** (1992) 192-197. Briefly, five ml of plasma are added to 96-well flat-bottom microtiter plates (Falcon), followed by the addition

of 5 ml of 80 mM MgCl_2 in incubation buffer (25 mM HEPES, 140 mM NaCl, 1 % RIA-grade BSA, pH 7.8). After a 5 min incubation at room temperature, the reaction is initiated by the addition of 10 ml of incubation buffer containing 0.1 mM substrate (H-Glycine-Proline-AMC; AMC is 7-amino-4-methylcoumarin). The plates are covered with aluminum foil (or kept in the dark) and incubated at room temperature for 20 min. After the 20 min reaction, fluorescence is measured using a CytoFluor 2350 fluorimeter (Excitation 380 nm Emission 460 nm; sensitivity setting 4). Test compounds are typically added as 2 ml additions and the assay buffer volume is reduced to 13 ml. A fluorescence-concentration curve of free AMC is generated using 0-50 mM solutions of AMC in assay buffer. The curve generated is linear and is used for interpolation of substrate consumption (catalytic activity in nmoles substrate cleaved/min). As with the previous assay, the potency of the test compounds as DPP-IV inhibitors, expressed as IC_{50} , is calculated from 8-point, dose-response curves using a 4 parameter logistic function.

In the above assay, IC_{50} values of from about 7 nM to about 2000 nM are obtained in human plasma, and of from about 3 nM to about 400 nM in rat plasma, e.g., for the agent of Example 3, of 7 nM in human and 6 nM in rat plasma, respectively.

In view of their ability to **inhibit DPP-IV**, the agents of the invention are indicated for use in **treating conditions mediated by DPP-IV**. It is expected that the compounds disclosed herein are useful in the treatment of **non-insulin-dependent diabetes mellitus, arthritis, obesity, and osteoporosis such as calcitonin-osteoporosis**. The agents of the invention improve early insulin response to an oral glucose challenge and, therefore, are particularly indicated for use in treating **non-insulin-dependent diabetes mellitus and further conditions of impaired glucose tolerance (IGT)**.

The ability of the agents of the invention to improve early insulin response to an oral **glucose challenge** may e.g. be measured in insulin resistant rats according to the following method:

Male Sprague-Dawley rats that have been fed a high fat diet (saturated fat = 57 % calories) for 2-3 weeks are fasted for approximately 2 hours on the day of testing, divided into groups of 8-10, and dosed orally with 10 mmol/kg of the test compounds in carboxymethylcellulose (CMC). An oral glucose bolus of 1 g/kg is administered 30 min after

the test compound directly into the stomach of the test animals. Blood samples, obtained at various timepoints from chronic jugular vein catheters are analyzed for plasma glucose and immunoreactive insulin (IRI) concentrations, and plasma DPP-IV activity. Plasma insulin levels are assayed by a double antibody radioimmunoassay (RIA) method using a specific anti-rat insulin antibody from Linco Research (St. Louis, MO, USA). The RIA has a lower limit of detection of 0.5 mU/ml with intra- and inter-assay variations of less than 5%. Data are expressed as % increase of the mean of the control animals.

It was found that upon oral administration, each of the compounds tested amplified the early insulin response which led to an improvement in glucose tolerance in the insulin resistant test animals. The following results were obtained:

Compound	Increase of insulin response at 10 mmol/kg
Ex. 1	61 %
Ex. 3	66 %
Ex. 5	108 %
Ex. 8	144 %
Ex. 12	59 %

The precise dosage of the agents of the invention to be employed for treating conditions mediated by DPP-IV inhibition depends upon several factors, including the host, the nature and the severity of the condition being treated, the mode of administration and the particular compound employed. However, in general, conditions mediated by DPP-IV inhibition are effectively treated when an agent of the invention is administered enterally, e.g. orally, or parenterally, e.g. intravenously, preferably orally, at a daily dosage of from about 0.002 mg/kg to about 5 mg/kg, preferably of from about 0.02 mg/kg to about 2.5 mg/kg body weight or, for most larger primates, a daily dosage of from about 0.1 mg to about 250 mg, preferably from about 1 mg to about 100 mg. A typical oral dosage unit is from about 0.01 mg/kg to about 0.75 mg/kg, one to three times a day. Usually, a small dose is administered initially and the dosage is gradually increased until the optimal dosage for the host

under treatment is determined. The upper limit of dosage is that imposed by side effects and can be determined by trial for the host being treated.

The agents of the invention may be combined with one or more pharmaceutically acceptable carriers and, optionally, one or more other conventional pharmaceutical adjuvants and administered enterally, e.g. orally, in the form of tablets, capsules, caplets, etc., or parenterally, e.g. intravenously, in the form of sterile injectable solutions or suspensions. The enteral and parenteral compositions may be prepared by conventional means.

The agents of the invention may be formulated into enteral and parenteral pharmaceutical compositions containing an amount of the active substance that is effective for treating conditions mediated by DPP-IV, such compositions in unit dosage form and such compositions comprising a pharmaceutically acceptable carrier.


Those agents of the invention which are e.g. of formula I may be administered in enantiomerically pure (S) form (e.g. $\geq 98\%$, preferably $\geq 99\%$ pure) or together with the other enantiomer, e.g. in racemic form. The above dosage ranges are based on the compounds of formula I (excluding the amount of R enantiomer).

The invention thus also comprises **an agent of the invention**, in particular, a compound of formula I as defined above, in free form or in pharmaceutically acceptable acid addition salt form, **for use as a pharmaceutical**. It further includes a **pharmaceutical composition** comprising an agent of the invention, in particular a compound of formula I as defined above, in free form or in pharmaceutically acceptable acid addition salt form, together with at least one pharmaceutically acceptable carrier or diluent. It further comprises the use of an agent of the invention, in particular, a compound of formula I as defined above, in free form or in pharmaceutically acceptable acid addition salt form **in the preparation of a medicament** for inhibiting DPP-IV or treating conditions mediated by DPP-IV, by a process comprising mixing an agent of the invention with a pharmaceutically acceptable carrier or diluent. It further provides a **method of inhibiting DPP-IV, or of treating conditions mediated by DPP-IV**, which comprises administering to a patient in need of such treatment a therapeutically effective amount of a compound of the invention, in particular, of formula I in free form or in pharmaceutically acceptable acid addition salt form.

The agents of Examples 1, 3, 5, 8 and 12 are the preferred agents of the invention, particularly those of Examples 1, 3, 5 and 12, preferably in hydrochloride acid addition salt form, especially the agent of Example 3, namely 1-[2-[(5-cyanopyridin-2-yl)amino]-ethylamino]acetyl-2-cyano-(S)-pyrrolidine, preferably in dihydrochloride acid addition salt form. It has been determined that in hydrochloride form they have an IC_{50} value in the Caco-2 DPP-IV assay of, respectively, 36, 22, 26, 8 and 279 nM, and in the modified Kubota assay above, an IC_{50} value for, respectively, human and rat plasma DPP-IV, of 27 and 22 nM (Example 1); 7 and 6 nM (Example 3); 37 and 18 nM (Example 5); 12 and 11 nM (Example 8); and 95 and 38 nM (Example 12). It is, therefore, indicated that for the above uses the compounds of Examples 1, 3, 5, 8 and 12 may be administered to larger mammals, for example humans, by similar modes of administration at similar dosages than conventionally employed with metformin.

Claims:

1. An N-(N'-substituted glycy)-2-cyanopyrrolidine.
2. A compound of formula I:



wherein **R** is:

- a) $R_1R_{1a}N(CH_2)_m$ wherein

R_1 is a pyridinyl or pyrimidinyl moiety optionally mono- or independently disubstituted with (C_{1-4}) alkyl, (C_{1-4}) alkoxy, halogen, trifluoromethyl, cyano or nitro; or phenyl optionally mono- or independently disubstituted with (C_{1-4}) alkyl, (C_{1-4}) alkoxy or halogen;

R_{1a} is hydrogen or (C_{1-8}) alkyl; and

m is 2 or 3;

- b) (C_{3-12}) cycloalkyl optionally monosubstituted in the 1-position with (C_{1-3}) hydroxyalkyl;

- c) $R_2(CH_2)_n$ wherein either

R_2 is phenyl optionally mono- or independently di- or independently trisubstituted with (C_{1-4}) alkyl, (C_{1-4}) alkoxy, halogen or phenylthio optionally monosubstituted in the phenyl ring with hydroxymethyl; or is (C_{1-8}) alkyl; a [3.1.1]bicyclic carbocyclic moiety optionally mono- or plurisubstituted with (C_{1-8}) alkyl; a pyridinyl or naphthyl moiety optionally mono- or independently disubstituted with (C_{1-4}) alkyl, (C_{1-4}) alkoxy or halogen; cyclohexene; or adamantyl; and

n is 1 to 3; or

R_2 is phenoxy optionally mono- or independently disubstituted with (C_{1-4}) alkyl, (C_{1-4}) alkoxy or halogen; and

n is 2 or 3;

- d) $(R_3)_2CH(CH_2)_2$ wherein each R_3 independently is phenyl optionally mono- or independently disubstituted with (C_{1-4}) alkyl, (C_{1-4}) alkoxy or halogen;

- e) $R_4(CH_2)_p$ - wherein R_4 is 2-oxopyrrolidinyl or (C_{2-4}) alkoxy and p is 2 to 4;
 - f) **isopropyl** optionally monosubstituted in 1-position with (C_{1-3}) hydroxyalkyl;
 - g) R_5 wherein R_5 is: indanyl; a pyrrolidinyl or piperidinyl moiety optionally substituted with benzyl; a [2.2.1]- or [3.1.1]bicyclic carbocyclic moiety optionally mono- or plurisubstituted with (C_{1-8}) alkyl; adamantyl; or (C_{1-8}) alkyl optionally mono- or independently plurisubstituted with hydroxy, hydroxymethyl or phenyl optionally mono- or independently disubstituted with (C_{1-4}) alkyl, (C_{1-4}) alkoxy or halogen;
- in free form or in acid addition salt form,

3. A compound according to claim 2 (a **compound Ip**) wherein R is R^P , which is:

- a) $R_1^P NH(CH_2)_2$ - wherein R_1^P is a pyridinyl or pyrimidinyl moiety optionally mono- or independently disubstituted with halogen, trifluoromethyl, cyano or nitro;
 - b) (C_{3-7}) cycloalkyl optionally monosubstituted in 1-position with (C_{1-3}) hydroxyalkyl;
 - c) $R_2^P(CH_2)_2$ - wherein R_2^P is phenyl optionally mono- or independently di- or independently trisubstituted with halogen or (C_{1-3}) alkoxy;
 - d) $(R_3^P)_2CH(CH_2)_2$ - wherein each R_3^P independently is phenyl optionally monosubstituted with halogen or (C_{1-3}) alkoxy;
 - e) $R_4(CH_2)_3$ - wherein R_4 is as defined above; or
 - f) **isopropyl** optionally monosubstituted in 1-position with (C_{1-3}) hydroxyalkyl;
- in free form or in acid addition salt form.

4. A compound according to claim 2 (a **compound Is**), wherein R is R^S , which is:

- a) $R_1^S R_{1a}^S (CH_2)_{ms}$ - wherein R_1^S is pyridinyl optionally mono- or independently disubstituted with chlorine, trifluoromethyl, cyano or nitro; pyrimidinyl optionally monosubstituted with chlorine or trifluoromethyl; or phenyl;
- R_{1a}^S is hydrogen or methyl; and
- ms is 2 or 3;
- b) (C_{3-12}) cycloalkyl optionally monosubstituted in 1-position with hydroxymethyl;

c) $R_2^s(CH_2)_{ns}$ - wherein either

R_2^s is phenyl optionally mono- or independently di- or independently trisubstituted with halogen, alkoxy of 1 or 2 carbon atoms or phenylthio monosubstituted in the phenyl ring with hydroxymethyl; (C_{1-6}) alkyl; 6,6-dimethylbicyclo[3.1.1]hept-2-yl; pyridinyl; naphthyl; cyclohexene; or adamantyl; and ns is 1 to 3; or R_2^s is phenoxy; and ns is 2;

d) (3,3-diphenyl)propyl;

e) $R_4^s(CH_2)_{ps}$ wherein R_4^s is 2-oxopyrrolidin-1-yl or isopropoxy and ps is 2 or 3;

f) isopropyl optionally monosubstituted in 1-position with hydroxymethyl;

g) R_5^s wherein R_5^s is: indanyl; a pyrrolidinyl or piperidinyl moiety optionally N-substituted with benzyl; bicyclo[2.2.1]hept-2-yl; 2,6,6-trimethylbicyclo[3.1.1]hept-3-yl; adamantyl; or (C_{1-8}) alkyl optionally mono- or independently disubstituted with hydroxy, hydroxymethyl or phenyl; in free form or in acid addition salt form.

5. The compound according to claim 2 wherein R is 2-[(5-cyanopyridin-2-yl)amino]ethyl, i.e. 1-[2-[(5-cyanopyridin-2-yl)amino]ethylamino]acetyl-2-cyano-(S)-pyrrolidine, in free form or in acid addition salt form, especially in dihydrochloride acid addition salt form, or a compound according to claim 2 which is of formula I wherein R is either


- 2-[(5-chloropyridin-2-yl)amino]ethyl, or
- (1-hydroxymethyl)cyclopent-1-yl, or
- 2-[(5-nitropyridin-2-yl)amino]ethyl, or
- 3-(isopropoxy)propyl,

in free form or in acid addition salt form.

6. A process for the preparation of a compound according to claim 1 which comprises coupling a reactive (2-cyanopyrrolidino)carbonylmethylene compound with an appropriate substituted amine or, for the preparation of a compound according to claim 2, which comprises

-25-

reacting a compound of formula II



wherein X is a reactive group,

with a compound of formula III



wherein R is as defined in claim 2,

and recovering the resultant compound in free form or in acid addition salt form.

7. A pharmaceutical composition comprising a compound according to claim 1 in free form or in pharmaceutically acceptable acid addition salt form, together with at least one pharmaceutically acceptable carrier or diluent.

8. A compound according to claim 1 in free form or in pharmaceutically acceptable acid addition salt form, for use as a pharmaceutical.

9. Use of a compound according to claim 1 in free form or in pharmaceutically acceptable acid addition salt form in the preparation of a medicament for inhibiting DPP-IV or treating conditions mediated by DPP-IV.

10. A method of inhibiting DPP-IV, or of treating a condition mediated by DPP-IV, which comprises administering a therapeutically effective amount of a compound according to claim 1 in free form or in pharmaceutically acceptable acid addition salt form to a patient in need of such treatment.

L6 ANSWER 3 OF 3 CAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 1998:323235 CAPLUS <<LOGINID::20071023>>
 DOCUMENT NUMBER: 129:16059
 TITLE: Preparation of N-aminoacetyl-2-cyanopyrrolidines as
 dipeptidylpeptidase IV inhibitors.
 INVENTOR(S): Villhauer, Edwin Bernard
 PATENT ASSIGNEE(S): Novartis A.-G., Switz.; Villhauer, Edwin Bernard
 SOURCE: PCT Int. Appl., 27 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9819998	A2	19980514	WO 1997-EP6125	19971105
WO 9819998	A3	19980716		
W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, SZ, BE, FR, GR, IE, IT, MC, NL, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
RW: GH, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
TW 492957	B	20020701	TW 1997-86115673	19971023
IN 1997MA02502	A	20050304	IN 1997-MA2502	19971103
CA 2269810	A1	19980514	CA 1997-2269810	19971105
CA 2269810	C	20060711		
AU 9853184	A	19980529	AU 1998-53184	19971105
AU 726186	B2	20001102		
EP 937040	A2	19990825	EP 1997-950112	19971105
EP 937040	B1	20040721		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, PT, IE, SI, FI, RO				
CN 1236361	A	19991124	CN 1997-199497	19971105
CN 1134411	B	20040114		
BR 9714130	A	20000229	BR 1997-14130	19971105
HU 2000000323	A2	20000828	HU 2000-323	19971105
HU 2000000323	A3	20020228		
JP 2000511559	T	20000905	JP 1998-521061	19971105
JP 3217380	B2	20011009		
RU 2180901	C2	20020327	RU 1999-111492	19971105
SK 282904	B6	20030109	SK 1999-608	19971105
AT 271543	T	20040815	AT 1997-950112	19971105
PT 937040	T	20041130	PT 1997-950112	19971105
ES 2226004	T3	20050316	ES 1997-950112	19971105
PL 190498	B1	20051230	PL 1997-332777	19971105
CZ 296303	B6	20060215	CZ 1999-1615	19971105
ZA 9709985	A	19980507	ZA 1997-9985	19971106
NO 9902028	A	19990428	NO 1999-2028	19990428
NO 312667	B1	20020617		
KR 2000053081	A	20000825	KR 1999-704004	19990506

HK 1022149
PRIORITY APPLN. INFO.:


A1 20050422

HK 2000-101080
US 1996-746295
WO 1997-EP6125

20000223
A 19961107
W 19971105

OTHER SOURCE(S):
GI

MARPAT 129:16059



AB Title compds. [I; R = (substituted) pyridinylaminoalkyl, pyrimidinylaminoalkyl, cycloalkyl, phenylalkyl, bicycloalkylalkyl, pyridinylalkyl, naphthylalkyl, phenoxyalkyl, etc.], were prepared. Thus, 2-[(5-chloropyridin-2-yl)amino]ethylamine in THF was treated with (S)-(2-cyanopyrrolidino)carbonylmethyl bromide in THF under ice cooling to give (S)-1-[2-[(5-chloropyridin-2-yl)amino]ethylamino]acetyl-2-cyanopyrrolidine dihydrochloride. I at 10 mmol/kg orally gave 59-144% increase in insulin response to an oral glucose challenge in rats.

IT 207612-01-9P


RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of N-aminoacetyl-2-cyanopyrrolidines as dipeptidylpeptidase IV inhibitors)

RN 207612-01-9 CAPLUS

CN 2-Pyrrolidinecarbonitrile, 1-[(bicyclo[2.2.1]hept-2-ylamino)acetyl]-, monohydrochloride, (2S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.



● HCl